



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

DNA methylation-based classification and grading system for meningioma: a multicentre, retrospective analysis

Sahm, Felix ; Schrimpf, Daniel ; Stichel, Damian ; Jones, David T W ; Hielscher, Thomas ; Schefzyk, Sebastian ; Okonechnikov, Konstantin ; Koelsche, Christian ; Reuss, David E ; Capper, David ; Sturm, Dominik ; Wirsching, Hans-Georg ; Berghoff, Anna Sophie ; Baumgarten, Peter ; Kratz, Annekathrin ; Huang, Kristin ; Wefers, Annika K ; Hovestadt, Volker ; Sill, Martin ; Ellis, Hayley P ; Kurian, Kathreena M ; Okuducu, Ali Fuat ; Jungk, Christine ; Drueschler, Katharina ; Schick, Matthias ; Bewerunge-Hudler, Melanie ; Mawrin, Christian ; Seiz-Rosenhagen, Marcel ; Ketter, Ralf ; Simon, Matthias ; et al

DOI: [https://doi.org/10.1016/S1470-2045\(17\)30155-9](https://doi.org/10.1016/S1470-2045(17)30155-9)

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-141060>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Sahm, Felix; Schrimpf, Daniel; Stichel, Damian; Jones, David T W; Hielscher, Thomas; Schefzyk, Sebastian; Okonechnikov, Konstantin; Koelsche, Christian; Reuss, David E; Capper, David; Sturm, Dominik; Wirsching, Hans-Georg; Berghoff, Anna Sophie; Baumgarten, Peter; Kratz, Annekathrin; Huang, Kristin; Wefers, Annika K; Hovestadt, Volker; Sill, Martin; Ellis, Hayley P; Kurian, Kathreena M; Okuducu, Ali Fuat; Jungk, Christine; Drueschler, Katharina; Schick, Matthias; Bewerunge-Hudler, Melanie; Mawrin, Christian; Seiz-Rosenhagen, Marcel; Ketter, Ralf; Simon, Matthias; et al (2017). DNA methylation-based classification and grading system for meningioma: a multicentre, retrospective analysis. *Lancet Oncology*, 18(5):682-694.

DOI: [https://doi.org/10.1016/S1470-2045\(17\)30155-9](https://doi.org/10.1016/S1470-2045(17)30155-9)

Molecular analysis of meningioma increases prognostic power: A methylation-based classification and grading system

Felix Sahm, MD^{1,2}, Daniel Schrimpf, PhD¹, Damian Stichel, PhD², David T.W. Jones, PhD³, Thomas Hielscher, MSc⁴, Sebastian Schefzyk, MSc¹, Konstantin Okonechnikov, PhD³, Christian Koelsche, MD^{1,2}, David Reuss, MD^{1,2}, David Capper, MD^{1,2}, Dominik Sturm, MD^{3,5}, Hans-Georg Wirsching, MD⁶, Anna Sophie Berghoff, MD⁷, Peter Baumgarten, MD⁸, Annekathrin Kratz, MD^{1,2}, Kristin Huang, MD^{1,2}, Annika K Wefers, MD^{1,2}, Volker Hovestadt, PhD⁹, Martin Sill, PhD⁴, Hayley P Ellis, BSc¹⁰, Kathreena M Kurian, MD¹⁰, Ali Fuat Okuducu, MD¹¹, Christine Jungk, MD¹², Katharina Drueschler, MD¹³, Matthias Schick¹⁴, Melanie Bewerunge-Hudler, PhD¹⁴, Christian Mawrin, MD¹⁵, Marcel Seiz-Rosenhagen, MD¹⁶, Ralf Ketter, MD¹⁷, Matthias Simon, MD¹⁸, Manfred Westphal¹⁹, MD, Katrin Lamszus, MD¹⁹, Albert Becker, MD²⁰, Arend Koch, MD²¹, Jens Schittenhelm, MD²², Elisabeth J Rushing²³, V Peter Collins, MD²⁴, Stefanie Brehmer, MD¹⁶, Lukas Chavez, PhD³, Michael Platten, MD^{13,25,26}, Daniel Hänggi, MD¹⁶, Andreas Unterberg, MD¹², Werner Paulus, MD²⁸, Wolfgang Wick, MD^{13,28}, Stefan M. Pfister, MD^{3,5}, Michel Mittelbronn, MD⁸, Matthias Preusser, MD²⁹, Christel Herold-Mende, PhD¹², Michael Weller, MD⁶, Andreas von Deimling, MD^{1,2}

1. Department of Neuropathology, Institute of Pathology, Ruprecht-Karls-University Heidelberg, Heidelberg, Germany,
2. Clinical Cooperation Unit Neuropathology, German Consortium for Translational Cancer Research (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany
3. Division of Pediatric Neurooncology, German Consortium for Translational Cancer Research (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany
4. Division of Biostatistics (C060), German Cancer Research Center (DKFZ), and German Cancer Consortium (DKTK), Heidelberg, Germany
5. Department of Pediatric Oncology, Haematology and Immunology, Heidelberg University Hospital, and National Center for Tumor Diseases (NCT), Heidelberg, Germany
6. Department of Neurology, University Hospital and University of Zurich, Zurich, Switzerland
7. Institute of Neurology, Comprehensive Cancer Center, CNS Tumours Unit (CCC-CNS), Medical University of Vienna, Austria
8. Neurological Institute (Edinger-Institute), Goethe University, Frankfurt, and German Cancer Consortium (DKTK), Germany
9. Division of Molecular Genetics, German Cancer Research Center (DKFZ), Heidelberg, Germany; current address: Broad Institute, Cambridge, MA, USA
10. Brain Tumour Research Group, Institute of Clinical Neurosciences, Southmead Hospital, University of Bristol, UK
11. Department of Pathology, University Hospital Nürnberg, Nürnberg, Germany
12. Department of Neurosurgery, University Hospital Heidelberg, Heidelberg, Germany
13. Neurology Clinic, Heidelberg University Hospital, Heidelberg, Germany
14. Genomics and Proteomics Core Facility, Micro-Array Unit, German Cancer Research Center, Heidelberg, Germany
15. Department of Neuropathology, Otto von Guericke University Magdeburg, Magdeburg, Germany
16. Department of Neurosurgery, University Hospital Mannheim, Mannheim, Germany
17. Department of Neurosurgery, Saarland University, Homburg, Germany
18. Department of Neurosurgery, Ev. Krankenhaus Bielefeld, Bielefeld, Germany
19. Department of Neurosurgery, University Hospital Hamburg-Eppendorf, Hamburg, Germany
20. Department of Neuropathology, University of Bonn, Bonn, Germany
21. Department of Neuropathology, Charité Medical University, Berlin, Germany
22. Department of Neuropathology, University Hospital Tübingen, Tübingen, Germany

- ^{23.} Department of Neuropathology, University Hospital and University of Zurich, Zurich, Switzerland
- ^{24.} Department of Molecular Histopathology, University of Cambridge, UK
- ^{25.} Clinical Cooperation Unit Neuroimmunology and Brain Tumor Immunology, German Consortium for Translational Cancer Research (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany
- ^{26.} Current address: Neurology Clinic, University Hospital Mannheim, Mannheim, Germany
- ^{27.} Institute of Neuropathology, University Hospital Münster, Münster, Germany
- ^{28.} Clinical Cooperation Unit Neurooncology, German Consortium for Translational Cancer Research (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany
- ^{29.} Department of Medicine I, Comprehensive Cancer Center, CNS Tumours Unit (CCC-CNS), Medical University of Vienna, Austria

Corresponding author

Dr. med. Felix Sahm and Prof. Dr. med. Andreas von Deimling

Department of Neuropathology

Institute of Pathology

Ruprecht-Karls-University Heidelberg, Heidelberg

Im Neuenheimer Feld 224

D-69120 Heidelberg

email: andreas.vondeimling@med.uni-heidelberg.de

phone: +49 6221 56 4651

Summary

Background

The World Health Organization (WHO) classification of brain tumors describes 15 subtypes of meningioma. Nine of these are allotted to WHO grade I, and three each to grade II and grade III, respectively. Grading is purely based on histology, with molecular markers lacking. While the current classification and grading approach is of prognostic value, it harbors shortcomings such as ill-defined parameters for subtypes and grading criteria prone to arbitrary judgment.

Methods

We investigated genome-wide DNA methylation patterns of 479 meningiomas to identify distinct methylation classes (MC) of meningioma. The MCs were further characterized by DNA copy-number analysis, mutational profiling and RNA sequencing. We validated our findings in an independent cohort of 140 tumors.

Findings

DNA methylation profiling distinguished six distinct MCs associated with typical mutational, cytogenetic, and gene expression patterns. Meningioma MCs exhibit a more homogeneous clinical course and allow prognostication with significantly higher power than the current morphology-based WHO classification. Meningioma MCs more accurately identify patients at high risk of recurrence among tumors with WHO grade I histology, and patients at lower risk of recurrence among WHO grade II tumors. DNA methylation-based classification and grading reduces the number of meningioma subtypes from 15, as historically defined by histology, to six clinically relevant MCs, each with a characteristic molecular and/or clinical profile.

Interpretation

DNA methylation-based meningioma classification captures biologically more homogenous groups and has a higher power for predicting tumor recurrence than the current WHO classification. The approach presented here is highly useful for stratifying meningioma patients for observation or adjuvant treatment groups. We consider methylation-based tumor classification highly relevant for the future diagnosis and treatment of meningioma.

Funding

This work was supported by the German Cancer Aid (110670, 110983) and the Else Kröner-Fresenius Foundation (A_60). We thank the DKFZ-Heidelberg for funding by HIPO H033.

Research in context

Evidence before this study

Meningiomas, the most frequent primary intracranial tumors, are diagnosed and graded according to the WHO classification of brain tumors. The recent update of this classification in 2016 has implemented molecular markers for several brain tumor entities.

However, there are still no established prognostic molecular markers for meningioma. Meningioma diagnostics is still based on purely histological criteria which are prone to a high inter-observer and sampling bias. Thus, the relevance of the current grading system for clinical decision making is heavily debated.

Previous work by several groups, including ours, has shown that DNA methylation signatures are specific for tumor entities. Importantly, DNA methylation profiling can identify biologically and clinically relevant subgroups among histologically indiscernible cases. Here, we employed this concept for the classification of meningiomas. A search in PubMed on October 21 2016 did not identify articles which used high-resolution DNA methylation profiling for identification of clinically relevant subgroups across all subtypes and grades of meningioma.

Added value of this study

We demonstrate that classification of meningiomas based on DNA methylation profiling is more powerful in predicting the clinical behavior than the current WHO classification and grading system. Our findings on a discovery series were confirmed on an independent validation series. Most notably, the novel approach was capable of identifying patients at high risk of rapid recurrence which were expected to have benign tumors based on WHO grading. Likewise, a considerable fraction of patients with the histological diagnosis of a higher grade meningioma - fostering the consideration of adjuvant treatment - but no recurrence could upfront be identified as low risk by DNA methylation profiling.

Implications of all the available evidence

Our data demonstrate that meningioma patients can be more accurately stratified for tumor behavior by DNA methylation profiling than by the current WHO classification. This greatly improves the basis for clinical decision making for or against additional therapy after surgery. We expect epigenetic profiling to be included into the diagnostic routine and implemented into upcoming updates of the WHO classification for brain tumors.

Introduction

The meninges exert a protective function for the entire central nervous system (CNS). During development, their precursor cells emerge from mesodermal structures and the neural crest, actively contributing to the differentiation of the brain¹⁻³. However, meningeal cells may transform to initiate tumors. These meningiomas are the most frequent primary intra-cranial and spinal tumors⁴. While 80 % of meningiomas show a benign clinical behavior and can be cured by resection alone, about 20 % recur and need additional treatment such as repeated surgery, irradiation, and systemic chemotherapy^{4,5}. Histopathological evaluation aims at the identification of cases at risk for recurrence. The histological differentiation into subtypes initially dates back to the 19th century. Later, in a first internationally recognized classification approach in 1928, Bailey and Cushing distinguished meningothelial, fibroblastic, and angiomatous subtypes⁶, and to this day, allocation to subtype is based solely on histological findings. The current WHO classification recognizes 15 subtypes and three grades of malignancy⁴, but some of the diagnostic criteria are vaguely defined and subject to a high inter-observer bias, indicating the need for more reliable markers^{5,7}.

For various other CNS tumors, molecular profiling has identified distinct subtypes with characteristic aberrations. Many of these correlate with prognosis or provide targets for treatment, and therefore support clinical decision making, e.g. epigenetic subgroups in medulloblastoma⁸⁻¹⁰ and ependymoma¹¹, or *isocitrate dehydrogenase (IDH)* status in diffuse glioma¹²⁻¹⁴. Recent studies identified *telomerase reverse transcriptase (TERT)* promoter mutations in a small subset of meningiomas to be associated with higher risk of recurrence and shorter time to progression^{15,16}, and four large exome-sequencing efforts focusing on WHO grade I meningiomas have identified recurrently mutated genes beyond the long-known association with *NF2*¹⁷⁻²⁰. Yet, these findings cover only a fraction of meningiomas and have not all been thoroughly tested for their prognostic relevance. In this study, we aimed at a comprehensive characterization of the entire molecular genetic landscape of meningioma in order to identify biologically and clinically relevant subgroups that refine the current classification scheme.

Results

DNA methylation analysis identifies six distinct methylation classes of meningioma

We generated genome-wide DNA methylation profiles from a discovery cohort of 497 meningiomas (Suppl. Fig. 1) along with 309 samples of other extra-axial skull tumors that may histologically mimic meningioma variants, including solitary fibrous tumor/hemangiopericytoma, schwannoma, malignant peripheral nerve sheath tumors, chordoma, chondrosarcoma, fibrous dysplasia, and hemangioblastoma. Despite sharing a mesodermal origin, unsupervised clustering of DNA methylation data clearly segregated all meningiomas from other skull tumors (Suppl. Fig 2). Unsupervised clustering of meningiomas alone revealed two major epigenetic groups (Groups A and B, Fig. 1A), with both groups further subdividing into four and two subgroups, respectively. These six subgroups were designated as “methylation classes” (MCs). Based on further molecular and clinical characteristics outlined below, the four MCs of Group A were designated MC benign 1 through 3 (MC ben-1, ben-2, ben-3) and MC intermediate A (MC int-A). The two MCs of Group B were designated MC intermediate B (MC int-B), and malignant (MC mal).

There was an enrichment of grade I tumors among MC ben-1, MC ben-2, and MC ben-3, and an enrichment of WHO grade III tumors in MC mal, while WHO grade II tumors were scattered across all MCs. Analysis of 75 primary and matched recurrent tumors from 37 patients showed that association with Group A or B was stable upon recurrence (Fig. 1B), supporting further assessment of methylation profiling for diagnostic and prognostic implications.

MC predict clinical course with higher accuracy than WHO grading

The wide spectrum of clinical behavior among WHO grade I and II meningiomas points towards the limited prognostic power of the current classification, particularly at the border between grade I and II. As a result, the basing of decisions about radiotherapy on the current grading scheme is heavily debated ⁵. Thus, we correlated meningioma MCs with progression-free survival (PFS) to evaluate their potential for predicting outcome compared to WHO grading (Fig. 2A, B). We further combined MCs exhibiting virtually identical benign (MC ben-1, MC ben-2, MC ben-3) or intermediate (MC int-A, MC int-B) outcome into combined MCs (Fig. 2C). Classification by individual and combined MCs demonstrates more precise prognostication than by WHO grading (Fig. 2D, Brier prediction test, $p < 0.01$). These findings were confirmed in 140 meningiomas from an independent validation cohort (Suppl. Fig 3A, B).

We next focused on the prediction power of MCs within WHO grades and, particularly, patients divergently diagnosed by WHO grading and DNA methylation-based classification. Patients with WHO grade I meningiomas who were molecularly assigned to an intermediate MC experienced a less

favorable clinical course than patients with WHO grade I meningiomas diagnosed solely based on histology. In fact, their outcome was indistinguishable from that of patients with WHO grade II meningiomas (Fig. 3A). Likewise, patients with WHO grade II meningiomas molecularly assigned to a benign MC had a better outcome than the average outcome of patients with histologically-defined WHO grade II meningiomas. Consequently, stratification for MC is of higher value for prediction of PFS than WHO grading. Within the combined MCs, WHO grading confers limited additional information (Fig. 3B, Suppl. Table 2). However, combined MCs delineate subgroups with significantly distinct prognosis within all WHO grades (Fig. 3C), demonstrating the benefit of MC-based grading for patients and the potential to significantly reduce under- or overtreatment.

Methylation classes are associated with distinct driver mutations and copy-number-alterations

We next sequenced 304 meningiomas with sufficient material available using a custom hybrid-capture next-generation sequencing (NGS) panel dedicated to 40 genes previously reported to be mutant in meningioma (Suppl. Table 1), based on our recently established custom NGS approach for routine brain tumor diagnostics²¹. Known recurrent mutations (most frequently *NF2*, followed by *TRAF7* and *AKT1*) were significantly enriched in certain MCs (Suppl. Table 3, Fig. 4). Within Group A, *NF2* mutations were observed in 63 % of MC ben-1 tumors (significant accumulation of parameter in this MC, $p < 0.0001$, Fisher's exact test). MC ben-2 contained the vast majority of meningiomas carrying *AKT1* (33 % in this subgroup; $p < 0.0001$), *SMO* (7 %; $p = 0.0002$), *KLF4* (15 %; $p < 0.0001$), and *TRAF7* (49 %; $p < 0.0001$) mutations, but only rarely harbored *NF2* mutations. Only one *AKT1* and five *KLF4* mutations were detected outside of MC ben-2. MC ben-3 exhibited *NF2* mutations in 32 % and *PIK3CA* mutations in 11 % of tumors, representing the majority (5/7, 71 %) of *PIK3CA* mutations in the cohort. MC int-A carried *NF2* mutations in 53 %. Within Group B, MC int-B tumors harbored *NF2* mutations in 35 % and MC mal in 31 % of cases. *SUFU* mutations were confined to Group B, with 5 % of MC int-B and 6 % MC mal tumors being mutated. Four out of five *TERT* promoter mutations mapped to meningiomas in Group B ($p = 0.005$, Fisher's exact test).

Annotation of copy-number-variations (CNV) revealed that MCs are associated with distinct cytogenetic aberrations (Fig. 4A, B): MC ben-1 was associated with deletions of 22q (95 %) but otherwise virtually no CNV. MC ben-2 presented with absence of recurrent CNVs. Typical for MC ben-3 were multiple chromosomal gains, most frequently affecting chromosome 5 (47 %). MC int-A frequently exhibited losses on 1p (70 %) and 22q (84 %). In Group B, MC int-B frequently exhibited losses on 1p (89 %), 10 and 22 (89 %), all features also shared with MC malignant. However, in MC mal, a higher frequency of *CDKN2A* deletion was apparent (70%).

Representative cases with sufficient material available of all MCs underwent RNA-sequencing which identified differentially upregulated genes and pathways (Suppl. Fig. 4).

Methylation classes and WHO subtypes, localization, and gender

Examining the distribution of histological subtypes, which currently determine grading, across MCs revealed which histological subtypes the MCs are composed of and, conversely, to which MC the samples of a respective subtype are assigned (Fig. 5). The rare lymphoplasmacyte-rich meningioma (WHO grade I) was not assessed due to the overwhelming dominance of constitutional (non-tumor) DNA in these samples. In general, two patterns were observed: Either a given MC was strongly associated with a small set of or even single histological subtypes, or samples of a particular MC or subtype were widely spread across all corresponding variants. MC ben-1 comprised the majority of fibroblastic meningiomas and is also enriched for psammomatous meningioma. Fibroblastic meningiomas frequently harbor calcifications called psammoma bodies, and a high abundance of these calcifications defines psammomatous meningioma. Their molecular similarity implies that they may rather represent a continuous spectrum of one phenotypic pattern. The overexpression of SERPINF1 (Suppl. Fig. 4), which has been implicated in osteogenesis and calcification, in MC ben-1 might contribute to this histologically detected phenomenon. MC ben-2 was highly enriched for meningothelial meningiomas, and contained the vast majority of secretory meningioma. MC ben-3 harbored cases from several subtypes but was particularly enriched for angiomatous meningiomas. This is in line with the overexpression of vessel-associated markers, e.g. factor VIII, in samples of this MC (Suppl. Fig. 4). Transitional meningioma, a hybrid of meningothelial and fibroblastic histology, dissolved into several MCs, along with the samples of the rare microcystic, chordoid, metaplastic, psammomatous, and clear cell subtype.

The two intermediate MCs were predominantly constituted of atypical meningiomas. However, a considerable fraction of atypical meningioma (n=31) fell into MC ben-1. The improved resolution of this group substantially contributes to the higher prognostic power of MC class over histology. Anaplastic meningiomas predominantly mapped to MC malignant. Of note, the six rhabdoid/papillary meningiomas, by definition WHO grade III, all ended up in one of the MC benign or intermediate meningioma groups. However, the number was too low to assess the statistical relevance of WHO grading and MC classification individually for rhabdoid/papillary meningioma.

Transitional meningioma WHO grade I was much more frequently assigned to an intermediate MC than fibroblastic or meningothelial meningioma, mostly to MC int-A. Atypical meningiomas assigned to a benign MC accumulated in MC ben-1.

The most frequent localizations for all subgroups were the frontal and central convexity, except for MC ben-2 (Fig 6). For the latter, basal localization was common, in line with the high occurrence of *AKT1* and *SMO* mutations in this MC which are known to be enriched in this location ^{19,22}. Interestingly, all MC mal cases were located at the convexity. In contrast, none of the basal tumors

were allotted to MC mal, including four intraventricular and ten spinal meningiomas that were all assigned to intermediate or benign MCs. Age distribution was equal throughout all MCs. In terms of gender, we observed a predominance of male patients in MC mal, while all other MCs mainly comprised female patients (Fig 6).

Discussion

The 15 subtypes of meningioma included in the current WHO classification have evolved over decades. The major aim of introducing this variety of subgroups was to cover the whole histological spectrum of meningioma and to avoid misclassification of tumors mimicking other entities. For example, meningeal tumors with chordoid or rhabdoid cytology may initially raise suspicion of a chordoma or rhabdoid tumor but not point towards meningioma. Therefore, particular subtypes with these features were introduced into the classification in order to draw attention to the morphologic diversity of meningioma. In addition, some cytological features have been reported to be associated with distinct outcome. Although this was based on small series, it prompted allotment of distinct WHO grades to specific meningioma subtypes. However, this approach has been increasingly questioned due to suboptimal inter-observer reproducibility and limited prognostic effect of the histological criteria of higher grade ^{7,23,24}, most recently in a large Radiation Therapy Oncology Group (RTOG) meningioma trial ⁷ in which the authors expressed the urgent need for more objective, molecular stratification markers.

This resulted in an overall critical view of clinicians with respect to the current meningioma WHO classification and grading, which has been expressed in the most recent published European Association of Neuro-oncology (EANO) guideline for the diagnosis and treatment of meningiomas⁵. Accordingly, revisiting meningioma diagnostics based on DNA methylation profiling by defining MCs with enhanced predictive power will greatly improve the acceptance of meningioma classification and more successfully guide decisions regarding postoperative treatment. An overview on the molecular and clinical hallmarks of the six meningioma MCs is given in Figure 6.

Distinct methylation profiles suggest different development

Beyond the identification of clinically relevant groups and the basis for a novel classification, our dataset might provide insight into the development of meningioma. This has previously been shown for other entities: Four variants of medulloblastoma, distinguishable by their DNA methylation patterns, were shown to arise from different precursor cell populations^{8,25-27}, and exhibit very different clinical characteristics and therapy needs. Our data indicate that the spectrum of meningiomas is divided into two major molecularly highly distinct Groups (A and B, Fig. 1). This strong separation suggests either the existence of distinct cells of origin or an underlying event with major impact on genome-wide DNA methylation. The very different DNA methylation profiles of Groups A and B despite the shared occurrence of *NF2* mutations might suggest that meningiomas arise from two different precursor cell populations. Based on our own and published high-throughput sequencing data, there is no evidence for the existence of a single mutation being solely responsible for the separation of these two groups. However, we cannot fully exclude the existence

of alterations not readily detectable by these approaches, such as translocations or fusions, causing the responsible changes in the methylome. Moreover, the fact that patients with meningiomas clustering in Group A share a predominantly benign, with a small proportion exhibiting an intermediate clinical course, and that patients with meningiomas of Group B follow an intermediate to malignant clinical course, may further argue towards a distinct cell of origin with different intrinsic propensities for malignant transformation. However, analyses dissecting the full regulatory background of the tumor cells in comparison to arachnoidal cells, e.g. by H3K27Ac ChIP-Sequencing, are needed to further elucidate this.

Methylation-based versus WHO subgrouping versus other molecular markers

Extensive whole exome or -genome sequencing has provided a large body of information on the mutational landscape of meningioma¹⁷⁻²⁰. Four distinct meningioma mutational subgroups have been proposed, defined by mutations either in *NF2*, *TRAF7*, the hedgehog pathway, or *POLR2A*¹⁸. However, such a model of meningioma development based on mutational analysis alone currently does not satisfy the clinical need for distinction between patients in need of adjuvant treatment or not. A major drawback is the lack of risk stratification among *NF2*-mutant cases, which can present with any clinical course. While the strong association of *AKT1*, *TRAF7/KLF4*, or *SMO* mutations with benign, or *TERT* promoter mutations with unfavorable course may allow for mutation-based risk assessment in these subgroups, the current inability to stratify *NF2*-mutated meningiomas for other mutational events associated with clinical outcome is a major obstacle for a classification and grading system based on mutational profiling alone.

Similarly strong limitations apply to approaches based on copy-number-profiles: They leverage the accumulation of aberrations during progression but are not capable of predicting the behavior upfront. The current dataset attributes the highest prognostic power to methylome-based subgrouping, which proves to be superior to WHO classification (Fig. 2, 3), while an exclusively mutation-based subgrouping for the full spectrum of meningioma is not available.

An integrated diagnosis for meningioma evaluation

The WHO 2016 revision of the classification for CNS tumors supports the concept of an integrated diagnosis. It relies on a multilayered approach combining data from histology, molecular genetic analyses, and clinical findings^{4,28,29}. Adopting this WHO approach to the diagnosis of meningioma, the morphological layer corresponds to the current diagnostic standard, i.e. diagnosing the 15 WHO meningioma subtypes and grading according to the morphological scheme. In the absence of molecular analyses the morphological diagnosis should be suffixed with NOS (not otherwise

specified), as agreed for parenchymal brain tumors without molecular workup⁴. The molecular diagnostic layer may contain elements such as DNA methylation and/or mutation analyses. Mutational data may enable inferring the MC for a subset within the MC ben-2, e.g. for *AKT1* mutant cases, but not in every instance. With methylation analysis performed, one of the six MCs can be diagnosed. If the MC is identified, this results in a significantly more powerful prediction of the clinical course. This corresponds to the current approach in other entities, e.g. ependymoma and medulloblastoma, for which methylation profiling has proven to be more relevant than histological grading^{4,11}. Based on the data presented here, the integrated diagnosis of meningioma will also highlight the prognostic impact of MCs, but in addition refer to the morphological subtype identified in histological examination.

Collectively, the dataset and accompanying classification scheme proposed here advances meningioma diagnostics from histology into an integrated profiling with higher accuracy of risk assessment for individual patients.

Author contributions

FS and AvD conceived the project, coordinated data generation, and wrote the manuscript with input from all co-authors. FS, D Schrimpf, D Stichel, DTWJ, SS, D Sturm, M Sill, VH, LC and SMP designed methylation experiments and analyzed DNA methylation data. FS, D Schrimpf, D Stichel, LC and KO analyzed RNA sequencing data. FS, D Schrimpf and TH analyzed survival data. FS, D Schrimpf, D Stichel, SS, DR, CK, DC, KH, AK, AW analyzed DNA sequencing data. M Schick and MBH performed array experiments. FS, DR, CK, DC, KH, AK, AW, PB, K Kurian, AFO, CM, AK, JS, EJ, VPC, WP, MM, and AvD performed histological evaluation. HGW, ASB, PB, HE, K Kurian, AFO, CM, CJ, KD, MSR, RK, M Simon, AB, M Westphal, KL, AK, JS, VPC, SB, M Platten, DH, AU, WP, WW, MM, M Preusser, CHM, and M Weller collected and interpreted clinical data and/or compiled respective tissue collections.

Acknowledgments

The authors thank Hai Yen Nguyen, Laura Doerner, Jochen Meyer and Julian Baron for excellent technical assistance. We also thank the Microarray unit of the Genomics and Proteomics Core Facility, German Cancer Research Center (DKFZ), especially Roger Fischer, Nadja Wermke and Anja Schramm-Glück, for providing excellent methylation services.

Materials and Methods

Samples

Samples with clinical data were retrospectively collected from the Dept. of Neuropathology Heidelberg, Germany (local and referral cases), Dept. of Neurosurgery Heidelberg and the FORAMEN network, the Dept. of Neurology and Neuropathology, Zürich, Switzerland, and the Neurological Institute (Edinger Institute) Frankfurt/Main, Germany. Additional samples without survival annotation were included from the Dept. of Neuropathology and Neurosurgery Berlin, Bonn, Hamburg, Magdeburg, Münster, Tübingen (all Germany), and Bristol (UK). The validation cohort was provided by the Medical University of Vienna. Sample and data collection was in accordance with local ethical regulations.

Methylation analysis, copy-number analysis

Unsupervised clustering of the 450k data of the discovery and the EPIC 850k data of the validation cohort was performed as previously described (see Supplementary Materials and Methods for references) based on Euclidian distance and Ward's linkage method. For the clustering probes with a standard deviation greater 0.2 across all samples were selected. The methylation probes were reordered by hierarchical clustering using Euclidean distance and complete linkage.

Copy-number aberrations were inferred from methylation array data using the R/Bioconductor package *conumee*.

Cohort-wide copy number analysis in MCs

Methylation-class wide relative copy-number assessment was performed based on 450k data by a proprietary algorithm and controlled by manual inspection of the *conumee*-based copy-number-profiles (Stichel et al., in preparation).

Panel and RNA sequencing

Panel sequencing for genes reported to be mutant in meningioma (Suppl. Table 1) was performed applying a custom hybrid-capture approach (Agilent) as described before. RNA libraries were generated with TruSeq RNA Access (Illumina) applying manufacturer supplied protocols. Sequencing was performed on a NextSeq 500 (Illumina).

Statistical analysis of clinical parameters

Distribution of survival times was estimated by the method of Kaplan and Meier and compared between groups with the log-rank test. Hazard ratios including 95% confidence intervals based on Cox regression models were calculated. For the multivariable Cox regression model, imputations of missing covariate values was done applying the multivariate imputations using chained equations

(mice) algorithm with 100 imputation runs. Hazard ratio for age is given per 10 year increment. Prediction error curves based on the Brier score were computed. Integrated Brier score was tested between risk stratifications using 1000 bootstrap samples. P-values below 0.05 were considered statistically significant. Analyses were performed with statistical software R 3.3.

Legends

Figure 1 Unsupervised clustering of methylation data of 479 meningioma samples (A). Unsupervised clustering of matched primary and recurrent samples (matched primary/recurrent samples of identical patient identified by arrows) combined with reference samples from group A and B shows that no shift between groups occurs upon recurrence (B).

Figure 2 Progression free survival (PFS) of 228 case with clinical data stratified for WHO grade (A), methylation class (B), combined methylation classes (C). Brier prediction plot calculated for the models A-C (D, WHO vs combined MCs $p=0.0138$, 0.0096 , 0.0062 for 5, 10 and 12 years, respectively).

Figure 3 Comparison of WHO grading and methylation-based risk prediction: WHO grade I cases allotted to an intermediate methylation class show PFS similar to the average grade II tumors. In turn, WHO grade II cases assigned to a benign methylation class have longer PFS than the average WHO grade II cases (A). Hazard ratio (including 95% confidence intervals) forest plot for WHO grading, overall and stratified for combined methylation classes (B). Hazard ratio forest plot for combined methylation classes, overall and stratified for WHO grading (C). While sub-stratification for WHO grade among MCs is of limited additional value (B), MCs stratify for distinct PFS within WHO grades (C).

Figure 4 Distribution of mutations across sample that underwent panel-sequencing (304) stratified for MCs (A). Copy number variations across all samples that underwent 450k analysis (497) the MCs (B).

Figure 5 Association of histological subtypes and MCs.

Figure 6 Schematic overview over the six identified MCs and their molecular and clinical characteristics

Supplementary Materials (online)

Supplementary Figure 1: Composition of Discovery Cohort (WHO grade, subtype, age, localization, gender)

Supplementary Figure 2: Unsupervised clustering of 450k methylation data of mesenchymal skull tumors and meningioma

Supplementary Figure 3: Kaplan-Meier analysis for Validation cohort stratified for WHO grade (A) and Methylation Class (B)

Supplementary Figure 4: Expression analysis based on RNA-sequencing data. Most differentially expressed genes in the six MCs (A) and ClueGo based on KEGG source data all samples (B). Nodes represent enriched gene sets, which are grouped and annotated by their similarity. Size is proportional to the number of involved genes. Manual curation was performed to remove uninformative sub-networks.

Supplementary Materials and Methods: Detailed methods with references

Supplementary Table 1: Multi-variate analyses

Supplementary Table 2: Genes included in panel sequencing

Supplementary Table 3: Distribution of mutations in Methylation classes


Supplementary Table 4: Clinical and mutational data of all cases

References


1. Mack J, Squier W, Eastman JT. Anatomy and development of the meninges: implications for subdural collections and CSF circulation. *Pediatr Radiol* 2009; **39**(3): 200-10.
2. Siegenthaler JA, Pleasure SJ. We have got you 'covered': how the meninges control brain development. *Curr Opin Genet Dev* 2011; **21**(3): 249-55.
3. Bifari F, Berton V, Pino A, et al. Meninges harbor cells expressing neural precursor markers during development and adulthood. *Front Cell Neurosci* 2015; **9**: 383.
4. Louis DN, Perry A, Reifenberger G, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol* 2016; **131**(6): 803-20.
5. Goldbrunner R, Minniti G, Preusser M, et al. EANO guidelines for the diagnosis and treatment of meningiomas. *Lancet Oncol* 2016; **17**(9): e383-91.
6. Bailey P, Cushing H. Angioblastic meningiomas. *Archives of pathology & laboratory medicine* 1928; **6**.
7. Rogers CL, Perry A, Pugh S, et al. Pathology concordance levels for meningioma classification and grading in NRG Oncology RTOG Trial 0539. *Neuro-oncology* 2016; **18**(4): 565-74.
8. Kool M, Korshunov A, Remke M, et al. Molecular subgroups of medulloblastoma: an international meta-analysis of transcriptome, genetic aberrations, and clinical data of WNT, SHH, Group 3, and Group 4 medulloblastomas. *Acta Neuropathol* 2012; **123**(4): 473-84.
9. Hovestadt V, Remke M, Kool M, et al. Robust molecular subgrouping and copy-number profiling of medulloblastoma from small amounts of archival tumour material using high-density DNA methylation arrays. *Acta Neuropathol* 2013; **125**(6): 913-6.
10. Zhukova N, Ramaswamy V, Remke M, et al. Subgroup-Specific Prognostic Implications of TP53 Mutation in Medulloblastoma. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2013; **31**(23): 2927-35.
11. Pajtler KW, Witt H, Sill M, et al. Molecular Classification of Ependymal Tumors across All CNS Compartments, Histopathological Grades, and Age Groups. *Cancer cell* 2015; **27**(5): 728-43.
12. Yan H, Parsons DW, Jin G, et al. IDH1 and IDH2 mutations in gliomas. *N Engl J Med* 2009; **360**(8): 765-73.
13. Hartmann C, Hentschel B, Wick W, et al. Patients with IDH1 wild type anaplastic astrocytomas exhibit worse prognosis than IDH1-mutated glioblastomas, and IDH1 mutation status accounts for the unfavorable prognostic effect of higher age: implications for classification of gliomas. *Acta Neuropathol* 2010; **120**(6): 707-18.
14. Cancer Genome Atlas Research N, Brat DJ, Verhaak RG, et al. Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *N Engl J Med* 2015; **372**(26): 2481-98.
15. Abedalthagafi MS, Bi WL, Merrill PH, et al. ARID1A and TERT promoter mutations in dedifferentiated meningioma. *Cancer Genet* 2015; **208**(6): 345-50.
16. Sahm F, Schrimpf D, Olar A, et al. TERT Promoter Mutations and Risk of Recurrence in Meningioma. *Journal of the National Cancer Institute* 2016; **108**(5).
17. Clark VE, Erson-Omay EZ, Serin A, et al. Genomic Analysis of Non-NF2 Meningiomas Reveals Mutations in TRAF7, KLF4, AKT1, and SMO. *Science* 2013.
18. Clark VE, Harmanci AS, Bai H, et al. Recurrent somatic mutations in POLR2A define a distinct subset of meningiomas. *Nature genetics* 2016.
19. Brastianos PK, Horowitz PM, Santagata S, et al. Genomic sequencing of meningiomas identifies oncogenic SMO and AKT1 mutations. *Nature genetics* 2013.
20. Reuss DE, Piro RM, Jones DT, et al. Secretory meningiomas are defined by combined KLF4 K409Q and TRAF7 mutations. *Acta Neuropathol* 2013.
21. Sahm F, Schrimpf D, Jones DT, et al. Next-generation sequencing in routine brain tumor diagnostics enables an integrated diagnosis and identifies actionable targets. *Acta Neuropathol* 2015.
22. Sahm F, Bissel J, Koelsche C, et al. AKT1E17K mutations cluster with meningothelial and transitional meningiomas and can be detected by SFRP1 immunohistochemistry. *Acta Neuropathol* 2013.



23. Vaubel RA, Chen SG, Raleigh DR, et al. Meningiomas With Rhabdoid Features Lacking Other Histologic Features of Malignancy: A Study of 44 Cases and Review of the Literature. *Journal of neuropathology and experimental neurology* 2016; **75**(1): 44-52.
24. Baumgarten P, Gessler F, Schittenhelm J, et al. Brain invasion in otherwise benign meningiomas does not predict tumor recurrence. *Acta Neuropathol* 2016; **132**(3): 479-81.
25. Lin CY, Erkek S, Tong Y, et al. Active medulloblastoma enhancers reveal subgroup-specific cellular origins. *Nature* 2016; **530**(7588): 57-62.
26. Northcott PA, Hielscher T, Dubuc A, et al. Pediatric and adult sonic hedgehog medulloblastomas are clinically and molecularly distinct. *Acta Neuropathol* 2011; **122**(2): 231-40.
27. Northcott PA, Shih DJ, Peacock J, et al. Subgroup-specific structural variation across 1,000 medulloblastoma genomes. *Nature* 2012; **488**(7409): 49-56.
28. Aldape K, Nejad R, Louis DN, Zadeh G. Integrating molecular markers into the World Health Organization classification of CNS tumors: a survey of the neuro-oncology community. *Neuro-oncology* 2016.
29. Louis DN, Perry A, Burger P, et al. International Society Of Neuropathology--Haarlem consensus guidelines for nervous system tumor classification and grading. *Brain pathology* 2014; **24**(5): 429-35.


A










B











Epigenetic and Biological Subgroups of Meningioma

